

References QBRI Insights: COVID-19 Treatment Recommendations – Latest Updates

1. Remdesivir (VEKLURY) [package insert]. Food and Drug Administration. 2020. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/214787Orig1s000lbl.pdf. Accessed: October 25, 2020
2. Beigel JH, Tomashek KM, Dodd LE, et al. Remdesivir for the treatment of COVID-19 - final report. *N Engl J Med*. 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32445440>.
3. Wang Y, Zhang D, Du G, et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet*. 2020;395(10236):1569-1578. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32423584>.
4. Food and Drug Administration. Fact sheet for healthcare providers: emergency use authorization (EUA) of casirivimab and imdevimab. 2020. Available at: <https://www.fda.gov/media/143892/download>. Accessed November 13, 2020.
5. Regeneron. Regeneron's COVID-19 outpatient trial prospectively demonstrates that REGN-COV2 antibody cocktail significantly reduced virus levels and need for further medical attention. 2020. Available at: <https://investor.regeneron.com/news-releases/news-release-details/regenerons-covid-19-outpatient-trial-prospectively-demonstrates>
6. Recovery Collaborative Group, Horby P, Lim WS, et al. Dexamethasone in hospitalized patients with COVID-19 - preliminary report. *N Engl J Med*. 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32678530>.
7. Tomazini BM, Maia IS, Cavalcanti AB, et al. Effect of dexamethasone on days alive and ventilator-free in patients with moderate or severe acute respiratory distress syndrome and COVID-19: the CoDEX randomized clinical trial. *JAMA*. 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32876695>.
8. Wang X, Guo X, Xin Q, et al. Neutralizing antibodies responses to SARS-CoV-2 in COVID-19 inpatients and convalescent patients. *Clin Infect Dis*. 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32497196>.
9. Gharbharan A, Jordans CCE, GeurtsvanKessel C, et al. Convalescent plasma for COVID-19: a randomized clinical trial. *medRxiv*. 2020;Preprint. Available at: <https://www.medrxiv.org/content/10.1101/2020.07.01.20139857v1>



10. Vincent MJ, Bergeron E, Benjannet S, et al. Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. *Virology*. 2005;2:69. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16115318>.
11. Liu J, Cao R, Xu M, et al. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discov*. 2020;6:16. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32194981>.
12. Maisonnasse P, Guedj J, Contreras V, et al. Hydroxychloroquine use against SARS-CoV-2 infection in non-human primates. *Nature*. 2020;585(7826):584-587.
13. Furtado RHM, Berwanger O, Fonseca HA, et al. Azithromycin in addition to standard of care versus standard of care alone in the treatment of patients admitted to the hospital with severe COVID-19 in Brazil (COALITION II): a randomised clinical trial. *Lancet*. 2020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32896292>.
14. Geleris J, Sun Y, Platt J, et al. Observational study of hydroxychloroquine in hospitalized patients with COVID-19. *N Engl J Med*. 2020;
15. Guaraldi G, Meschiari M, Cozzi-Lepri A, Milic J, Tonelli R, Menozzi M, Franceschini E, Cuomo G, Orlando G, Borghi V et al: Tocilizumabin patients with severe COVID-19: a retrospective cohort study. *The Lancet Rheumatology* 2020, 2(8):e474-e484
16. Somers EC, Eschenauer GA, Troost JP, Golob JL, Gandhi TN, Wang L, Zhou N, Petty LA, Baang JH, Dillman NO et al: Tocilizumab for treatment of mechanically ventilated patients with COVID-19. *medRxiv : the preprint server for health sciences* 2020.
17. Cantini F, Goletti D, Petrone L, Najafi Fard S, Niccoli L, Foti R. Immune Therapy, or Antiviral Therapy, or Both for COVID-19: A Systematic Review. *Drugs*. 2020 Dec;80(18):1929-1946. doi: 10.1007/s40265-020-01421-w. PMID: 33068263; PMCID: PMC7568461